

Diptera (flies) recorded at Holt Island Reserve, St. Ives (TL310714) during 2002

	17.6	17.7	18.8	13.9	7.10
Tipulidae					
<i>Nephrotoma quadrifaria</i> (Meigen)	/	/			
<i>Tipula lateralis</i> Meigen			/		
<i>paludosa</i> Meigen				/	
Limoniidae					
<i>Achyrolimonia decemmaculata</i> (Loew)				/	
<i>Austrolimnophila ochracea</i> (Meigen)		/			/
<i>Cheilotrichia cinerascens</i> (Meigen)	/				
<i>Dicranomyia chorea</i> (Meigen)	/				
<i>modesta</i> (Meigen)					/
<i>Erioptera fusculentata</i> Edwards					/
<i>Limonia nubeculosa</i> (Meigen)	/	/	/	/	/
<i>macrostigma</i> (Schummel)			/		/
<i>Neolimonia dumetorum</i> (Meigen)		/			
<i>Neolimnomyia nemoralis</i> (Meigen)					/
<i>Molophilus griseus</i> (Meigen)		/			
<i>Pilaria discicollis</i> (Meigen)					/
<i>Rhipidia maculata</i> (Meigen)				/	
Bibionidae					
<i>Dilophus febrilis</i> L.			/		
Mycetophilidae					
<i>Acnemia nitidicollis</i> (Meigen)				/	
<i>Cordyla crassicornis</i> Meigen					/
<i>Mycetophila curviseta</i> Lundstrom					/
<i>dentata</i> Lundstrom					/
<i>mitis</i> Johannsen				/	
<i>Mycomya prominens</i> (Lundstrom)			/		
<i>Sciophila nonnisilva</i> Hutson	/		/		
Sciaridae					
<i>Bradysia inusitata</i> Tuomikoski				/	
<i>Scaptosciara vitripennis</i> (Meigen)	/		/		
<i>Schwenkfeldina carbonaria</i> (Meigen)			/	/	
Psychodidae					
<i>Boreoclytocerus ocellaris</i> (Meigen)	/		/	/	
<i>Pericoma compta</i> Eaton					/
<i>nubila</i> (Meigen)		/			
<i>trivialis</i> Eaton					/
<i>Trichomyia urbica</i> Haliday	/				
Anisopodidae					
<i>Sylvicola cinctus</i> (Fabricius)	/			/	/
<i>punctatus</i> (Fabricius)	/				
Ptychopteridae					
<i>Ptychoptera albimana</i> (Fabricius)				/	
Ceratopogonidae					
<i>Clinohelea unimaculata</i> (Macquart)	/				
<i>Mallochohelea nitida</i> (Macquart)	/				
<i>setigera</i> (Loew)	/	/			
<i>Palpomyia luteifemorata</i> Edwards	/				
Rhagionidae					
<i>Chrysopilus asiliformis</i> (Preysslner)		/			

	17.6	17.7	18.8	13.9	7.10
Stratiomyidae					
Beris vallata (Forster)	/	/			
Chloromyia formosa (Scopoli)		/			
Microchrysa polita (L.)		/			
Pachygaster leachii Stephens		/			
Hybotidae					
Bicellaria vana Collin	/		/		
Drapetis ephippiata (Fallén)	/				
humilis Frey			/		
simulans Collin		/			
Ocydromia glabricula (Fallén)	/	/	/		/
Leptopeza flavipes (Meigen)			/		
Platypalpus annulipes (Meigen)	/				
calcaratus (Meigen)	/				
cothurnatus Macquart	/				
cursitans (Fabricius)	/				
longisetus Zetterstedt	/				
minutus (Meigen)	/			/	
nigritarsis (Fallén)				/	
pallidiventris (Meigen)		/			
Empididae					
Empis albinervis Meigen	/				
caudatula Loew	/				
livida L.	/				
Hemerodromia baetica Collin	/				
Hilara intermedia (Fallén) <i>interstincta</i> (Fallén)	/				
monedula Collin	/				
Rhamphomyia crassirostris (Fallén)	/				
nigripennis (Fabricius)	/	/			
Dolichopodidae					
Argyra leucocephala (Meigen)		/			/
Campsicnemus curvipes (Fallén)	/			/	/
Chrysotus gramineus (Fallén)	/	/			
Dolichopus brevipennis Meigen	/				
cilifemoratus Macquart	/				
festivus Haliday	/	/	/		
latelimbatus Macquart				/	
plumipes (Scopoli)	/	/		/	
ungulatus (L.)		/			
Hercostomus nanus (Macquart)	/				
Poecilobothrus nobilitatus (L.)	/		/		
Rhaphium appendiculatum Zetterstedt				/	
caliginosum Meigen		/	/	/	
Sympycnus desoutterii Parent				/	
Syntormon bicolorillum (Zetterstedt)			/		/
denticulatum (Zetterstedt)		/			
pallipes (Fabricius)		/			/
Teucophorus calcaratus (Macquart)	/	/	/		
nigricosta (von Roser)	/				
Opetiidae					
Opetia nigra Meigen					/

Phoridae

Phora atra (Meigen) /

Lonchopteridae

Lonchoptera bifurcata (Fallén) /

lutea Panzer / / / / /

Syrphidae

Cheilosia albitarsis (Meigen) /

Eupeodes luniger (Meigen) /

Episyrphus balteatus (De Geer) /

Eristalis pertinax (Scopoli) /

tenax (L.) / /

Melanostoma scalare (Fabricius) /

Myathropa florea (L.) / /

Platycheirus albimanus (Fabricius) / /

scutatus (Meigen) / /

Rhingia campestris (Meigen) /

Syrphus ribesii (L.) /

Volucella inanis (L.) H /

pellucens (L.) /

Micropezidae

Neria cibaria (L.) /

Psilidae

Chamaepsila pallida (Fallén) /

rosae (Fabricius) /

Chyliza leptogaster (Panzer) /

Conopidae

Conops quadrifasciatus De Geer /

Lonchaeidae

Dasiops spatiosus (Becker) /

Lonchaea chorea (Fabricius) /

mallochi MacGowan & Rotheray /

Palloppteridae

Palloptera muliebris (Harris) /

trimacula (Meigen) / / /

ustulata Fallén / / /

Ulidiidae

Melieria crassipennis (Fabricius) /

Seioptera vibrans (L.) /

Tephritidae

Anomoia permunda (Harris) /

Phylophylla caesio (Harris) / /

Lauxaniidae

Caliopum simillimum Collin / / /

Homoneura tesquae (Becker) /

thalhammari Papp /

Lyciella decempunctata (Fallén) / /

rorida (Fallén) / /

Minettia longipennis (Fabricius) / /

Sapromyza halidayi Shatalkin /

sempunctata Meigen /

Tricholauxania praeusta (Fallén) /

	17.6	17.7	18.8	13.9	7.10
Sciomyzidae					
<i>Limnia unguicornis</i> (Scopoli)		/			
<i>Pherbellia schonherri</i> (Fallén)		/			
<i>Sciomyza dryomyzina</i> Zetterstedt		/			
<i>Tetanocera arrogans</i> Meigen		/			
<i>hyalipennis</i> von Roser	/	/	/		/
Sepsidae					
<i>Nemopoda nitidula</i> (Fallén)		/			
<i>Sepsis cynipsea</i> (L.)			/		
<i>fulgens</i> Meigen	/		/		
<i>punctum</i> (Fabricius)	/				
Clusiidae					
<i>Clusiodes albimana</i> (Meigen)				/	/
<i>gentilis</i> (Collin)					/
Agromyzidae					
<i>Amauromyza flavifrons</i> (Meigen)				/	
<i>labiatarum</i> Hendel		H		/	
<i>Agromyza mobilis</i> Meigen		/			
<i>nigripes</i> Meigen	/				
<i>pseudoreptans</i> Nowakowski			/		
<i>Cerodontha muscina</i> (Meigen)		H	/		
<i>Chromatomyia milii</i> Kaltenbach	/	/			
<i>Melanagromyza angeliciphaga</i> Spencer		/			
<i>Phytomyza ranunculi</i> (Schrank)	/	/		/	
<i>tetrasticha</i> Hendel		/			
Opomyzidae					
<i>Geomyza tripunctata</i> Fallén	/				
<i>Opomyza florum</i> (Fabricius)		/	/	/	
<i>germinationis</i> (L.)	/	/	/	/	/
Anthomyzidae					
<i>Paranthomyza nitida</i> (Meigen)				/	
Asteidae					
<i>Asteia amoena</i> Meigen	/	/			
Milichiidae					
<i>Desmometopa sordida</i> Fallén)				/	
Carnidae					
<i>Meoneura vagans</i> Fallén		/			
Chloropidae					
<i>Calamoncosis aspistyliana</i> Duda	RDB K	H	/		
<i>glyceriae</i> Nartshuk			/		
<i>Cetema neglectum</i> Tonnoir			/		
<i>simile</i> Ismay <i>elongatum</i> (Meigen)			/		
<i>Chlorops hypostigma</i> Meigen	/		/		
<i>pumilionis</i> (Bjerkander)			/		/
<i>Dicraeus vagans</i> (Meigen)		/			
<i>Elachiptera cornuta</i> (Fallén)		/	/		/
<i>megaspis</i> (Loew)					/
<i>Oscinella nitidissima</i> (Meigen)	/				
<i>trochanterata</i> Collin	/				
<i>Thaumatomyia glabra</i> (Meigen)		/			
<i>notata</i> (Meigen)	/				/
<i>Tricimba cincta</i> (Meigen)	/				

	17.6	17.7	18.8	13.9	7.10
Heleomyzidae					
Heteromyza rotundicornis (Zetterstedt)	/				
Sphaeroceridae					
Ischiolepta nitida (Duda) H			/		
pusilla (Fallén)			/	/	
Leptocera fontinalis (Fallén)		/	/	/	
lutosa (Stenhammar)			/		/
Opacifrons coxata (Stenhammar)			/		
Pseudocollinella humida (Haliday)					/
Pullimosina moesta (Villeneuve)			/		
Spelobia ochripes (Meigen)					/
rufulabris (Stenhammar)			/		
Trachysiphonella leucoptera (Haliday) <i>Trachyopella</i>			/		
Drosophilidae					
Drosophila andalusiaca Strobl		/		/	/
confusa Staeger				/	
fenestrarum Fallén				/	
immigrans Sturtevant					/
melanogaster Meigen					/
obscura Fallén					/
subobscura Collin					/
Leucophenga maculata Dufour				/	
Scaptomyza pallida (Zetterstedt)	/	/	/	/	/
Ephydriidae					
Caenia palustris (Fallén)		/			
Ditrichophora calceata (Meigen)	/		/		
Gymnoclasiopa plumosa (Fallén)		/			
Limnellia quadrata (Fallén)	/			/	/
Notiphila riparia Meigen	/	/			
umbrosa Drake H	/				
Parydra coarctata (Fallén)	/	/	/	/	/
fossarum (Haliday)		/			
Pelina aenea (Fallén)	/				
Philygria interstincta (Fallén)		/			
vittipennis (Zetterstedt) H		/			
Scatella tenuicosta Collin				/	
Scathophagidae					
Cleigastra apicalis (Meigen)		/	/		
Cordilura ciliata Meigen		/			
Norellisoma spinimanum (Fallén)	/	/	/		
Scathophaga furcata (Say)	/			/	
stercoraria (L.)	/			/	
Anthomyiidae					
Anthomyia procellaris Rondani		/			
Botanophila fugax (Meigen)		/	/	/	
striolata (Fallén)		/			
Delia platura (Meigen)	/		/		
Hydrophoria lancifer (Harris)		/			
ruralis (Meigen)				/	
Hylemya vagans (Panzer)				/	/
Hylemyza partita (Meigen)		/	/		
Lasiomma seminitidum (Zetterstedt)				/	

	17.6	17.7	18.8	13.9	7.10
Pegomya flavifrons (Walker)				/	
ulmaria Rondani				/	
Pegoplata infirma (Meigen)	/				
Fanniidae					
Fannia armata (Meigen)	/	/		/	
coracina (Loew)				/	
fuscata (Fallén)	/				
manicata (Meigen)	/		/		
Muscidae					
Azelia cilipes (Haliday)				/	
nebulosa Robineau-Desvoidy				/	
zetterstedti Rondani	/				
Coenosia agromyzina (Fallén)	/	/	/		/
infantula Rondani		/	/		
Graphomya maculata (Scopoli)				/	
Hebecnema nigricolor (Fallén)			/		
vespertina (Fallén)		/	/	/	
Helina evecata (Harris)	/				
Hydrotaea armipes (Fallén)				/	
similis Meade			/		
Mesembrina meridiana (L.)			/		
Musca autumnalis De Geer			/		
Muscina levida (Harris)				/	/
Phaonia halterata (Stein)	/		/		
palpata (Stein)					/
subventa (Harris)	/	/	/	/	
valida (Harris)				/	
Calliphoridae					
Melanomyia nana (Meigen)	/	/	/		
Sarcophagidae					
Sarcophaga subvicina Rohdendorf			/		
Tachinidae					
Meigenia mutabilis (Fallén)	/	/	/		
Siphona geniculata (De Geer)	/				

Nomenclature follows Chandler, P.J. 1998 Checklists of Insects of the British Isles (New Series) Pt.1 : Diptera. *Handbk Ident. Br. Insects* 12. 234pp. Royal Entomological Society.

Red Data Book status follows Falk, S. 1991. A review of the scarce and threatened flies of Great Britain (Pt.1). Nature Conservancy Council, Peterborough. **RDB 2** = Vulnerable. **RDB K** = insufficiently known but suspected to fall within an RDB category.

H = new record for Huntingdonshire.

Dec. 2002.

Jonathan Cole
2, Lenton Close
Brampton
Huntingdon
Cambs PE28 4TR